

Princeton University Observatory
Peyton Hall
Princeton, NJ 08544

7/23/96
6:00
6/1/96

INTERIM REPORT

The Universe at Moderate Redshift

NASA Grant NAG 5-2759

Covering the Period
July 1, 1995 through June 30, 1996

Submitted to

National Aeronautics and Space Administration
Goddard Space Flight Center
Greenbelt, MD 20771

Prepared by: Renyue Cen
Research Astronomer

And: Jeremiah P. Ostriker
Principal Investigator

July 23, 1996

PROGRESS REPORT

for

NASA GRANT #NAG 5-2759

During the Period July 1, 1995 through June 30, 1996

Progress Report

The work done in the past year by J.P. Ostriker, R. Cen, J.R. Gott and collaborators covered a wide range of fields including origin of the cosmological X-ray background, properties of X-ray clusters, a comparison of the thermodynamic properties of a suite of currently popular cosmological models, topological properties of galaxy distributions as a function of galaxy type, nonlinear patterns in gravitational clustering process, development of a ray tracing algorithm to study the gravitational lensing by cosmic structures, effects of weak gravitational lensing by large-scale structure on the determination of q_0 , and implementation of a new accurate hydro-N-body code. We, in collaboration with other groups in our Grand Challenge Cosmology Consortium, are now undertaking a major study of the four most attractive cosmological models, including tilted $\Omega = 1$ cold dark matter model, low Ω cold dark matter model with a cosmological constant, mixed hot and cold matter model, and low Ω open cold dark matter model. We will study all models in a spectrum of scales from Lyman Alpha clouds to clusters of galaxies with unprecedented resolutions. This undertaking will enable us to seriously constrain/reject the models, by comparing the models' predictions with observations by HST, ROSAT and eventually AXAF. The details of the last year's studies are now described.

R. Cen, J.P. Ostriker and collaborators present a new treatment of two popular models for the growth of structure, examining the X-ray emission from hot gas with allowance for spectral line emission from various atomic species, primarily "metals". The X-ray emission from the bright cluster sources is not significantly changed from prior work and, as noted earlier, shows the CDM+ Λ model (LCDM) to be consistent but the standard, COBE normalized model (SCDM) to be inconsistent with existing observations — after allowance for still the considerable numerical modelling uncertainties. But they find one important new result. Radiation in the softer band 0.5-1.0keV is predominantly emitted by gas far from cluster centers (hence "background"). This background emission dominates over the cluster emission below 1keV and observations of it should show clear spectral signatures indicating its origin. In particular the "iron blend" should be seen prominently in this spectral bin from cosmic background hot gas at high galactic latitudes and should show shadowing against the SMC indicating its extragalactic origin. Certain OVII lines also provide a signature of this gas which emits a spectrum characteristic of $10^{6.6 \pm 0.6}$ K gas. Recent ASCA observations of the X-ray background tentatively indicate the presence of component with exactly the spectral features they predict there.

R. Cen, J.P. Ostriker and collaborators have compared the observed optical and X-ray properties of clusters of galaxies with hydrodynamic simulations of clusters in two cosmological models : standard ($\Omega = 1$) and low-density flat CDM models, normalized to the COBE background fluctuations. The observed properties include the velocity dispersion

versus temperature relation, the gas mass versus total mass relation, and the velocity dispersion versus gas mass fraction relation of clusters. They find that both models reproduce well the *shape* of the observed functions : $\sigma \sim T^{0.5}$, implying the gas is in approximate hydrostatic equilibrium with the cluster potential and an approximately constant ratio of gas to total mass in clusters. The *amplitude* of the relations, however, differs between the models : low-density CDM is consistent with the data, while $\Omega = 1$ CDM is not. The observed β parameter of clusters, $\beta = \sigma^2/(kT/\mu m_p) = 0.94 \pm 0.08$, is consistent with that produced by low-density CDM but considerably larger, by nearly a factor of two, than expected for $\Omega = 1$. Low-density CDM is also more consistent with the average observed relation of $M_{gas} \sim 0.16 M h_{50}^{-1.5}$ for clusters, while $\Omega = 1$ CDM yields a gas mass that is four times lower. The cluster gas mass fraction reflects approximately the baryon fraction in the models, Ω_b/Ω . Due to its larger baryon fraction, a low-density model is more consistent with the observed gas mass fraction in clusters than is an $\Omega = 1$ model.

J.P. Ostriker and R. Cen have computed, including a current state-of-the-art treatment of hydrodynamical processes, heating and cooling, a variety of cosmological models into the extreme nonlinear phase to enable comparisons with observations. The results for such a suite of currently interesting models are summarized and compared. First, we note the common, model independent results. All have a mean ($z = 0$) temperature of $10^{4.5} - 10^{5.5}$ K, set essentially by photoheating processes. Most gas is in one of two components: either at the photoheating floor $10^{4.5}$ K and primarily in low density regions or else shock heated to $10^5 - 10^6$ K and in regions of moderate overdensity (in caustics and near groups and clusters). It presents a major observational challenge to observationally detect this second, abundant component as it is neither an efficient radiator nor absorber. About 2% to 10% of the baryons cool and collapse into galaxies forming on caustics and migrating to clusters. About 1%-2% of baryons are in the very hot X-ray emitting gas near cluster cores, in good agreement with observations. These correspondances between the simulations and the real world imply that there is some significant truth to the underlying standard scenarios for the growth of structure. The differences among model predictions may help us find the path to the correct model. For COBE normalized models the most relevant differences concern epoch of structure formation. In the open variants having $\Omega = 0.3$, with or without a cosmological constant, structure formation on galactic scales is well advanced at redshift $z=5$, and reionization occurs early. But if observations require models for which most galaxy formation occurs more recently than $z = 2$, then the flat $\Omega = 1$ models are to be preferred. The velocity dispersion on the $1h^{-1}\text{Mpc}$ scale also provides a strong discriminant with, as expected, the $\Omega = 1$ models giving a much higher (perhaps too high) a value for that statistic.

J.R. Gott, R. Cen and J.P. Ostriker have studied the topology of large scale structure as a function of galaxy type using the genus statistic. In hydrodynamical cosmological CDM simulations, galaxies form on caustic surfaces (Zeldovich pancakes) then slowly drain onto filaments and clusters. The earliest forming galaxies in the simulations (defined as "ellipticals") are thus seen at the present epoch preferentially in clusters (tending toward a meatball topology), while the latest forming galaxies (defined as "spirals") are seen currently in a spongelike topology. The topology is measured by the genus (= number

of “donut” holes - number of isolated regions) of the smoothed density-contour surfaces. The measured genus curve for all galaxies as a function of density obeys approximately the theoretical curve expected for random-phase initial conditions, but the early forming elliptical galaxies show a shift toward a meatball topology relative to the late forming spirals. Simulations using standard biasing schemes fail to show such an effect. Large observational samples separated by galaxy type could be used to test for this effect.

J.P. Ostriker and collaborators successfully implemented the particle-particle/particle-mesh (P³M) code on the special purpose hardware GRAPE. The code currently achieves a peak efficiency of one-third the speed of a vectorized P³M code on a Cray C-90, and significant improvements are planned in the near future.

R. Cen, J.P. Ostriker and collaborators have investigated the nonlinear clustering of dark matter particles in an expanding universe using N-body simulations. One can gain some insight into this complex problem if simple relations between physical quantities in the linear and nonlinear regimes can be extracted from the results of N-body simulations. Hamilton et al. (1991) and Nityananda and Padmanabhan (1994) have made an attempt in this direction by relating the mean relative pair velocities to the mean correlation function in a useful manner. They investigate this relation and other closely related issues in detail for the case of six different power spectra: power laws with spectral indexes $n = -2, -1$, cold dark matter (CDM), and hot dark matter models with density parameter $\Omega = 1$; CDM including a cosmological constant (Λ) with $\Omega_{CDM} = 0.4$, $\Omega_{\Lambda} = 0.6$; and $n = -1$ model with $\Omega = 0.1$. They find that: (i) Power law spectra lead to self-similar evolution in an $\Omega = 1$ universe. (ii) Stable clustering does not hold in an $\Omega = 1$ universe to the extent our simulations can ascertain. (iii) Stable clustering is a better approximation in the case of $\Omega < 1$ universe in which structure formation freezes out at some low redshift. (iv) The relation between dimensionless pair velocity and the mean correlation function, $\bar{\xi}$, is only approximately independent of the shape of the power spectrum. At the nonlinear end, the asymptotic value of the dimensionless pair velocity decreases with increasing small scale power, because the stable clustering assumption is not universally true. (v) The relation between the evolved $\bar{\xi}$ and the linear regime $\bar{\xi}$ is also not universal but shows a weak spectrum dependence. They present simple theoretical arguments for these conclusions.

R. Cen, J.P. Ostriker and collaborators have described in detail a new method to trace light rays through an essentially three dimensional mass distribution up to high redshift. As an example, we apply this method here to a standard cold dark matter universe. We obtain a variety of results, some of them statistical in nature, others from rather detailed case studies of individual “lines of sight”. Among the former are the frequency of multiply imaged quasars, the distribution of separation of the multiple quasars, and the redshift distribution of lenses: all that as a function of quasar redshift. We find effects from very weak lensing up to highly magnified multiple images of high redshift objects. Applied to extended sources, i.e. galaxies, this ranges from slight deformations of the shapes, only measurable in a big ensemble, through tangentially aligned arclets up to giant luminous arcs. We can study the weak coherent shear fields produced by lensing of large scale structure in directions that are devoid of large mass concentrations as well as the strong

lensing around massive clusters of galaxies. Gravitational lensing directly measures mass density fluctuations along the line of sight to very distant objects. No assumptions need to be made concerning bias, the ratio of fluctuations in galaxy density to mass density. Hence lensing is a good tool to study the universe at medium and high redshifts. Cosmological models – normalized to the universe at redshift zero – differ considerably in their predictions for the mass distributions at these distance scales. Therefore lensing is a powerful tool to distinguish between various cosmological models. Our ultimate goal is to apply this method to a number of cosmogonic models in order to study their gravitational lensing effects and be able to eliminate some models whose properties are very different from the properties of the observed universe.

R. Cen, J.P. Ostriker and collaborators have examined effects of the weak gravitational lensing by large-scale structure on the determination of the cosmological deceleration parameter q_0 . They find that the lensing induced dispersions on truly standard candles are 0.04 and 0.02 mag at redshift $z = 1$ and $z = 0.5$, respectively, in a COBE-normalized cold dark matter universe with $\Omega_0 = 0.40$, $\Lambda_0 = 0.6$, $H = 65\text{km/s/Mpc}$ and $\sigma_8 = 0.79$. It is shown that one would observe $q_0 = -0.44^{+0.17}_{-0.05}$ and $q_0 = -0.45^{+0.10}_{-0.03}$ (the errorbars are 2σ limits) with standard candles with zero intrinsic dispersion at redshift $z = 1$ and $z = 0.5$, respectively, compared to the truth of $q_0 = -0.40$ in this case, i.e., a 10% error in q_0 will be made. A standard COBE normalized $\Omega_0 = 1$ CDM model would produce three times as much variance and a mixed (hot and cold) dark matter model would lead to an intermediate result. One unique signature of this dispersion effect is its non Gaussianity. Although the lensing induced dispersion at lower redshift is still significantly smaller than the currently best observed (total) dispersion of 0.12 mag in a sample of type Ia supernovae, selected with the multicolor light curve shape method, it becomes significant at higher redshift. They show that there is an optimal redshift, in the range $z \sim 0.5 - 2.0$ depending on the amplitude of the intrinsic dispersion of the standard candles, at which q_0 can be most accurately determined.

R. Cen and J.P. Ostriker, in collaboration with G.L. Bryan, M.L. Norman and J.M. Stone of University of Illinois, described a hybrid scheme for cosmological simulations that incorporates a Lagrangean particle-mesh (PM) algorithm to follow the collisionless matter with the higher order accurate piecewise parabolic method (PPM) to solve the equations of gas dynamics. Both components interact through the gravitational potential, which requires the solution of Poisson's equation, here done by Fourier transforms. Due to the vast range of conditions that occur in cosmological flows (pressure difference of up to fourteen orders of magnitude), a number of additions and modifications to PPM were required to produce accurate results. These are described, as are a suite of cosmological tests.

Publications Supported by NASA Grant #NAG 5-2759

During the Period July 1, 1995 through June 30, 1996

Papers Published:

1. "A Cosmological Hydrodynamic Code Based on the Piecewise Parabolic Method"
G.L. Bryan, M.L. Norman, J.M. Stone, R. Cen and J.P. Ostriker (1995).
The Computational Physics Journal, 89, 149
2. "Background X-ray Emission from Hot Gas in CDM and CDM+ Λ Universes: Spectral Signatures"
R. Cen, H. Kang, J.P. Ostriker and D. Ryu (1995)
Astrophysical Journal, 451, 436
3. "Cosmological Simulations Using Special Purpose Computers: Implementing P³M on GRAPE"
P.P. Brieu, F.J. Summers and J.P. Ostriker (1995)
Astrophysical Journal, 453, 566
4. "The Baryon Fraction and Velocity-Temperature Relation in Galaxy Clusters: Models versus Observations"
L.M. Lubin, R. Cen, N. A. Bahcall and J.P. Ostriker (1996)
Astrophysical Journal, 460, 10
5. "Hydrodynamic Simulations of the Growth of Cosmological Structure: Summary and Comparisons Among Scenarios"
J.P. Ostriker and R. Cen (1996)
Astrophysical Journal, 464, 27
6. "Topology of Large-scale Structure by Galaxy Type: Hydrodynamic Simulations"
J.R. Gott, R. Cen and J.P. Ostriker (1996)
Astrophysical Journal, 465, 499

Papers in press:

7. "Patterns in Nonlinear Gravitational Clustering: A Numerical Investigation"
T. Padmanabhan, R. Cen, J.P. Ostriker, and F J Summers (1996)
Astrophysical Journal, in press
8. "The Lyman Alpha Forest from Gravitational Collapse in the CDM+ Λ Model"
J. Miralda-Escude, R. Cen, J.P. Ostriker and M. Rauch (1996)
Astrophysical Journal, in press
9. "Effects of Weak Gravitational Lensing on the Determination of q_0 "
J. Wambsganss, R. Cen, G. Xu and J. P. Ostriker (1996)
Astrophysical Journal Letters, in press

Papers submitted:

10. "Testing Cosmological Models by Gravitational Lensing: I. Method and First Applications"
J. Wambsganss, R. Cen and J.P. Ostriker (1996)
Astrophysical Journal, submitted

PROGRESS REPORT

for

NASA GRANT #NAG 5-2759

During the Period July 1, 1995 through June 30, 1996

Progress Report

The work done in the past year by J.P. Ostriker, R. Cen, J.R. Gott and collaborators covered a wide range of fields including origin of the cosmological X-ray background, properties of X-ray clusters, a comparison of the thermodynamic properties of a suite of currently popular cosmological models, topological properties of galaxy distributions as a function of galaxy type, nonlinear patterns in gravitational clustering process, development of a ray tracing algorithm to study the gravitational lensing by cosmic structures, effects of weak gravitational lensing by large-scale structure on the determination of q_0 , and implementation of a new accurate hydro-N-body code. We, in collaboration with other groups in our Grand Challenge Cosmology Consortium, are now undertaking a major study of the four most attractive cosmological models, including tilted $\Omega = 1$ cold dark matter model, low Ω cold dark matter model with a cosmological constant, mixed hot and cold matter model, and low Ω open cold dark matter model. We will study all models in a spectrum of scales from Lyman Alpha clouds to clusters of galaxies with unprecedented resolutions. This undertaking will enable us to seriously constrain/reject the models, by comparing the models' predictions with observations by HST, ROSAT and eventually AXAF. The details of the last year's studies are now described.

R. Cen, J.P. Ostriker and collaborators present a new treatment of two popular models for the growth of structure, examining the X-ray emission from hot gas with allowance for spectral line emission from various atomic species, primarily "metals". The X-ray emission from the bright cluster sources is not significantly changed from prior work and, as noted earlier, shows the CDM+ Λ model (LCDM) to be consistent but the standard, COBE normalized model (SCDM) to be inconsistent with existing observations — after allowance for still the considerable numerical modelling uncertainties. But they find one important new result. Radiation in the softer band 0.5-1.0keV is predominantly emitted by gas far from cluster centers (hence "background"). This background emission dominates over the cluster emission below 1keV and observations of it should show clear spectral signatures indicating its origin. In particular the "iron blend" should be seen prominently in this spectral bin from cosmic background hot gas at high galactic latitudes and should show shadowing against the SMC indicating its extragalactic origin. Certain OVII lines also provide a signature of this gas which emits a spectrum characteristic of $10^{6.6 \pm 0.6} \text{K}$ gas. Recent ASCA observations of the X-ray background tentatively indicate the presence of component with exactly the spectral features they predict there.

R. Cen, J.P. Ostriker and collaborators have compared the observed optical and X-ray properties of clusters of galaxies with hydrodynamic simulations of clusters in two cosmological models : standard ($\Omega = 1$) and low-density flat CDM models, normalized to the COBE background fluctuations. The observed properties include the velocity dispersion

versus temperature relation, the gas mass versus total mass relation, and the velocity dispersion versus gas mass fraction relation of clusters. They find that both models reproduce well the *shape* of the observed functions : $\sigma \sim T^{0.5}$, implying the gas is in approximate hydrostatic equilibrium with the cluster potential and an approximately constant ratio of gas to total mass in clusters. The *amplitude* of the relations, however, differs between the models : low-density CDM is consistent with the data, while $\Omega = 1$ CDM is not. The observed β parameter of clusters, $\beta = \sigma^2/(kT/\mu m_p) = 0.94 \pm 0.08$, is consistent with that produced by low-density CDM but considerably larger, by nearly a factor of two, than expected for $\Omega = 1$. Low-density CDM is also more consistent with the average observed relation of $M_{gas} \sim 0.16 M h_{50}^{-1.5}$ for clusters, while $\Omega = 1$ CDM yields a gas mass that is four times lower. The cluster gas mass fraction reflects approximately the baryon fraction in the models, Ω_b/Ω . Due to its larger baryon fraction, a low-density model is more consistent with the observed gas mass fraction in clusters than is an $\Omega = 1$ model.

J.P. Ostriker and R. Cen have computed, including a current state-of-the-art treatment of hydrodynamical processes, heating and cooling, a variety of cosmological models into the extreme nonlinear phase to enable comparisons with observations. The results for such a suite of currently interesting models are summarized and compared. First, we note the common, model independent results. All have a mean ($z = 0$) temperature of $10^{4.5} - 10^{5.5}$ K, set essentially by photoheating processes. Most gas is in one of two components: either at the photoheating floor $10^{4.5}$ K and primarily in low density regions or else shock heated to $10^5 - 10^6$ K and in regions of moderate overdensity (in caustics and near groups and clusters). It presents a major observational challenge to observationally detect this second, abundant component as it is neither an efficient radiator nor absorber. About 2% to 10% of the baryons cool and collapse into galaxies forming on caustics and migrating to clusters. About 1%-2% of baryons are in the very hot X-ray emitting gas near cluster cores, in good agreement with observations. These correspondances between the simulations and the real world imply that there is some significant truth to the underlying standard scenarios for the growth of structure. The differences among model predictions may help us find the path to the correct model. For COBE normalized models the most relevant differences concern epoch of structure formation. In the open variants having $\Omega = 0.3$, with or without a cosmological constant, structure formation on galactic scales is well advanced at redshift $z=5$, and reionization occurs early. But if observations require models for which most galaxy formation occurs more recently than $z = 2$, then the flat $\Omega = 1$ models are to be preferred. The velocity dispersion on the $1h^{-1}\text{Mpc}$ scale also provides a strong discriminant with, as expected, the $\Omega = 1$ models giving a much higher (perhaps too high) a value for that statistic.

J.R. Gott, R. Cen and J.P. Ostriker have studied the topology of large scale structure as a function of galaxy type using the genus statistic. In hydrodynamical cosmological CDM simulations, galaxies form on caustic surfaces (Zeldovich pancakes) then slowly drain onto filaments and clusters. The earliest forming galaxies in the simulations (defined as “ellipticals”) are thus seen at the present epoch preferentially in clusters (tending toward a meatball topology), while the latest forming galaxies (defined as “spirals”) are seen currently in a spongelike topology. The topology is measured by the genus (= number

of “donut” holes - number of isolated regions) of the smoothed density-contour surfaces. The measured genus curve for all galaxies as a function of density obeys approximately the theoretical curve expected for random-phase initial conditions, but the early forming elliptical galaxies show a shift toward a meatball topology relative to the late forming spirals. Simulations using standard biasing schemes fail to show such an effect. Large observational samples separated by galaxy type could be used to test for this effect.

J.P. Ostriker and collaborators successfully implemented the particle-particle/particle-mesh (P³M) code on the special purpose hardware GRAPE. The code currently achieves a peak efficiency of one-third the speed of a vectorized P³M code on a Cray C-90, and significant improvements are planned in the near future.

R. Cen, J.P. Ostriker and collaborators have investigated the nonlinear clustering of dark matter particles in an expanding universe using N-body simulations. One can gain some insight into this complex problem if simple relations between physical quantities in the linear and nonlinear regimes can be extracted from the results of N-body simulations. Hamilton et al. (1991) and Nityananda and Padmanabhan (1994) have made an attempt in this direction by relating the mean relative pair velocities to the mean correlation function in a useful manner. They investigate this relation and other closely related issues in detail for the case of six different power spectra: power laws with spectral indexes $n = -2, -1$, cold dark matter (CDM), and hot dark matter models with density parameter $\Omega = 1$; CDM including a cosmological constant (Λ) with $\Omega_{CDM} = 0.4$, $\Omega_{\Lambda} = 0.6$; and $n = -1$ model with $\Omega = 0.1$. They find that: (i) Power law spectra lead to self-similar evolution in an $\Omega = 1$ universe. (ii) Stable clustering does not hold in an $\Omega = 1$ universe to the extent our simulations can ascertain. (iii) Stable clustering is a better approximation in the case of $\Omega < 1$ universe in which structure formation freezes out at some low redshift. (iv) The relation between dimensionless pair velocity and the mean correlation function, $\bar{\xi}$, is only approximately independent of the shape of the power spectrum. At the nonlinear end, the asymptotic value of the dimensionless pair velocity decreases with increasing small scale power, because the stable clustering assumption is not universally true. (v) The relation between the evolved $\bar{\xi}$ and the linear regime $\bar{\xi}$ is also not universal but shows a weak spectrum dependence. They present simple theoretical arguments for these conclusions.

R. Cen, J.P. Ostriker and collaborators have described in detail a new method to trace light rays through an essentially three dimensional mass distribution up to high redshift. As an example, we apply this method here to a standard cold dark matter universe. We obtain a variety of results, some of them statistical in nature, others from rather detailed case studies of individual “lines of sight”. Among the former are the frequency of multiply imaged quasars, the distribution of separation of the multiple quasars, and the redshift distribution of lenses: all that as a function of quasar redshift. We find effects from very weak lensing up to highly magnified multiple images of high redshift objects. Applied to extended sources, i.e. galaxies, this ranges from slight deformations of the shapes, only measurable in a big ensemble, through tangentially aligned arclets up to giant luminous arcs. We can study the weak coherent shear fields produced by lensing of large scale structure in directions that are devoid of large mass concentrations as well as the strong

lensing around massive clusters of galaxies. Gravitational lensing directly measures mass density fluctuations along the line of sight to very distant objects. No assumptions need to be made concerning bias, the ratio of fluctuations in galaxy density to mass density. Hence lensing is a good tool to study the universe at medium and high redshifts. Cosmological models – normalized to the universe at redshift zero – differ considerably in their predictions for the mass distributions at these distance scales. Therefore lensing is a powerful tool to distinguish between various cosmological models. Our ultimate goal is to apply this method to a number of cosmogonic models in order to study their gravitational lensing effects and be able to eliminate some models whose properties are very different from the properties of the observed universe.

R. Cen, J.P. Ostriker and collaborators have examined effects of the weak gravitational lensing by large-scale structure on the determination of the cosmological deceleration parameter q_0 . They find that the lensing induced dispersions on truly standard candles are 0.04 and 0.02 mag at redshift $z = 1$ and $z = 0.5$, respectively, in a COBE-normalized cold dark matter universe with $\Omega_0 = 0.40$, $\Lambda_0 = 0.6$, $H = 65\text{km/s/Mpc}$ and $\sigma_8 = 0.79$. It is shown that one would observe $q_0 = -0.44^{+0.17}_{-0.05}$ and $q_0 = -0.45^{+0.10}_{-0.03}$ (the errorbars are 2σ limits) with standard candles with zero intrinsic dispersion at redshift $z = 1$ and $z = 0.5$, respectively, compared to the truth of $q_0 = -0.40$ in this case, i.e., a 10% error in q_0 will be made. A standard COBE normalized $\Omega_0 = 1$ CDM model would produce three times as much variance and a mixed (hot and cold) dark matter model would lead to an intermediate result. One unique signature of this dispersion effect is its non Gaussianity. Although the lensing induced dispersion at lower redshift is still significantly smaller than the currently best observed (total) dispersion of 0.12 mag in a sample of type Ia supernovae, selected with the multicolor light curve shape method, it becomes significant at higher redshift. They show that there is an optimal redshift, in the range $z \sim 0.5 - 2.0$ depending on the amplitude of the intrinsic dispersion of the standard candles, at which q_0 can be most accurately determined.

R. Cen and J.P. Ostriker, in collaboration with G.L. Bryan, M.L. Norman and J.M. Stone of University of Illinois, described a hybrid scheme for cosmological simulations that incorporates a Lagrangean particle-mesh (PM) algorithm to follow the collisionless matter with the higher order accurate piecewise parabolic method (PPM) to solve the equations of gas dynamics. Both components interact through the gravitational potential, which requires the solution of Poisson's equation, here done by Fourier transforms. Due to the vast range of conditions that occur in cosmological flows (pressure difference of up to fourteen orders of magnitude), a number of additions and modifications to PPM were required to produce accurate results. These are described, as are a suite of cosmological tests.

Publications Supported by NASA Grant #NAG 5-2759

During the Period July 1, 1995 through June 30, 1996

Papers Published:

1. "A Cosmological Hydrodynamic Code Based on the Piecewise Parabolic Method"
G.L. Bryan, M.L. Norman, J.M. Stone, R. Cen and J.P. Ostriker (1995).
The Computational Physics Journal, 89, 149
2. "Background X-ray Emission from Hot Gas in CDM and CDM+ Λ Universes: Spectral Signatures"
R. Cen, H. Kang, J.P. Ostriker and D. Ryu (1995)
Astrophysical Journal, 451, 436
3. "Cosmological Simulations Using Special Purpose Computers: Implementing P³M on GRAPE"
P.P. Brieu, F.J. Summers and J.P. Ostriker (1995)
Astrophysical Journal, 453, 566
4. "The Baryon Fraction and Velocity-Temperature Relation in Galaxy Clusters: Models versus Observations"
L.M. Lubin, R. Cen, N. A. Bahcall and J.P. Ostriker (1996)
Astrophysical Journal, 460, 10
5. "Hydrodynamic Simulations of the Growth of Cosmological Structure: Summary and Comparisons Among Scenarios"
J.P. Ostriker and R. Cen (1996)
Astrophysical Journal, 464, 27
6. "Topology of Large-scale Structure by Galaxy Type: Hydrodynamic Simulations"
J.R. Gott, R. Cen and J.P. Ostriker (1996)
Astrophysical Journal, 465, 499

Papers in press:

7. "Patterns in Nonlinear Gravitational Clustering: A Numerical Investigation"
T. Padmanabhan, R. Cen, J.P. Ostriker, and F J Summers (1996)
Astrophysical Journal, in press
8. "The Lyman Alpha Forest from Gravitational Collapse in the CDM+ Λ Model"
J. Miralda-Escude, R. Cen, J.P. Ostriker and M. Rauch (1996)
Astrophysical Journal, in press
9. "Effects of Weak Gravitational Lensing on the Determination of q_0 "
J. Wambsganss, R. Cen, G. Xu and J. P. Ostriker (1996)
Astrophysical Journal Letters, in press

Papers submitted:

10. "Testing Cosmological Models by Gravitational Lensing: I. Method and First Applications"
J. Wambsganss, R. Cen and J.P. Ostriker (1996)
Astrophysical Journal, submitted